


# Procedure Card

## COVID-19 Antigen Rapid Test Device

### Step 2 - Option A: Nasal Swab (Anterior Nares)



#### Step 2a.1

Tilt patient's head 70°. Insert the swab at least 0.5 inch inside the nostril (nares) until mild resistance was encountered at the middle turbinate.



#### Step 2a.2

Using a circular motion, the nasal orifice should be swabbed for a minimum of five seconds.



#### Step 2a.3

Compress the nostril with the fingers to trap the swab tip and rotate the tip for a minimum of five seconds.



#### Step 2a.4


Repeat steps for other nostril.

### Step 2 - Option B: Nasopharyngeal Swab

#### Step 2b

Tilt patient's head 70°. Insert the swab into the nostril parallel to the palate, and gently push the swab into the posterior nasopharynx.

Rotate against nasal wall to ensure swab contains cells and mucus.




#### Step 3

Insert the swab into the extraction tube. Mix well and squeeze the swab 10-15 times. After waiting 2 minutes, twist the swab against the inner wall as you remove it, trying to release as much liquid as you can while removing.



#### Step 4

Insert the nozzle into the sample extraction tube.



#### Step 5


Add 3 drops of the solution into the sample well by gently squeezing the tube.



#### Step 6

Read results after 15 minutes.

### Result Interpretations



#### Step 1

Gently mix the extraction buffer.

Add 10 drops into the extraction tube.

